

Radiative decays of charmonium into light mesons

Ying-Jia Gao, Yu-Jie Zhang, Kuang-Ta Chao
Department of Physics, Peking University, Beijing 100871

We apply perturbative QCD to the radiative decays of charmonia J/ψ and χ_{cJ} into light mesons. We perform a complete numerical calculation for the quark-gluon loop diagrams involved in these processes. The calculated J/ψ decay branching ratios into P-wave mesons $f_2(1270)$ and $f_1(1285)$ fit the data well, while that of $f_0(980)$ (if treated as an $s\bar{s}$ meson) is predicted to be 1.6×10^{-4} , which implies that $f_0(1710)$ can not be the $s\bar{s}$ or $(u\bar{u} + d\bar{d})/\sqrt{2}$ meson. Decays of P-wave charmonia $\chi_{cJ} \rightarrow \rho(\omega, \phi)\gamma$ ($J=0,1,2$) are also studied, and the branching ratio of $\chi_{c1} \rightarrow \rho\gamma$ is predicted to be 1.4×10^{-5} , which may be tested by CLEO-c and BESIII with future experiments.

Decays of heavy quarkonium into light hadrons are forbidden processes by the OZI rules. The radiative decays of J/ψ into light hadrons such as $J/\psi \rightarrow \gamma f_2(1270)$ are expected to proceed via two virtual gluons that subsequently convert to light mesons, with the photon emitted by the charm quarks (see Fig.1 for the three Feynman diagrams). These processes are interesting since they can provide useful information for understanding the fundamental theory of strong interactions, namely, quantum chromodynamics (QCD).

In this letter, we will report calculations for radiative decays of charmonia J/ψ and χ_{cJ} into various light mesons based on nonrelativistic quantum chromodynamics. Hopefully, the J/ψ mass is large enough for using perturbative QCD to calculate these processes.

Differing from previous works (see, e.g., [1, 2]), we will perform a complete numerical calculation for the quark-gluon loop diagrams involved in these processes.

Here, it is emphasized that in our approach the description of J/ψ as a nonrelativistic $c\bar{c}$ bound state is a good approximation; while treating the light mesons such as $f_2(1270)$ as nonrelativistic $q\bar{q}$ bound states ($q = u, d, s$) with constituent quark masses $m_u = m_d \approx 350 \text{ MeV}$, $m_s \approx 500 \text{ MeV}$ is only an approximation for describing the nonperturbative dynamics that has been used in the nonrelativistic constituent quark models rather successfully, and was also used in previous studies for these radiative decay processes [1, 2]. In fact, if the current quark masses for the light quarks, which will vanish in the chiral limit, were used in the quark-gluon loop diagram, we would encounter difficulties with soft and collinear divergences.

In the following, we will first focus on the radiative decays of J/ψ to P-wave light mesons such as $f_2(1270)$, $f_1(1285)$, and $f_0(980)$. We adopt the descriptions of nonrelativistic bound states, i.e., both J/ψ and light mesons are described by the color-singlet non-relativistic wave functions. As widely accepted assignments we assume that $f_2(1270)$ and $f_1(1285)$ are mainly composed of $(u\bar{u} + d\bar{d})/\sqrt{2}$ (neglecting the mixing with $s\bar{s}$ for simplicity). But for $f_0(980)$, there are many possible assignments such as the tetraquark state, the $K\bar{K}$ molecule, and the P-wave $s\bar{s}$ dominated state (for related discussions on $f_0(980)$ and other scalar mesons, see, e.g., the topical review-note on scalar mesons in [3] and [4]). Since experimental data show that $D_s^+ \rightarrow f_0\pi^+$ has a large branching ratio (BR) [3], we prefer to assign $f_0(980)$ as an $s\bar{s}$ dominated P-wave state as a tentative choice.

In the calculations, the spin projection operators for $^3S_1$, $^1S_0$ and ^3P_J states and their relations can be found in [5, 6]. We use `FeynCalc` [7] for the tensor reduction and `LoopTools` [8] for the numerical evaluation of infrared safe integrals. We follow the way in Ref. [9] to deal with five-point functions and high tensor loop integrals that can not be calculated by `LoopTools` and `FeynCalc`, such as

$$E_{\alpha\beta\rho\sigma} = \int d^D k \frac{k_\alpha k_\beta k_\rho k_\sigma}{k^2 [(k + p_q)^2 - m_q^2] (k + 2p_q)^2 [(p_c + k)^2 - m_c^2] [(k + 2p_q - p_c)^2 - m_c^2]}, \quad (1)$$

where p_c is the momentum of c quark, p_q the momentum of light quark, m_c the charm quark mass, and m_q the constituent light quark mass of f_J meson. For the constituent quark masses, we choose $m_u = m_d = 0.35 \text{ GeV}$, $m_s = 0.50 \text{ GeV}$, $m_c = 1.5 \text{ GeV}$.

Using the theoretical expression for the widths of $f_2 \rightarrow \gamma\gamma$,

$$\Gamma_{f_2(1270) \rightarrow \gamma\gamma}^{(th)} = \frac{6N_c}{5} (Q_u^2 + Q_d^2)^2 \alpha^2 \frac{|\mathcal{R}'_P(0)|^2}{m^4}, \quad \Gamma_{f'_2(1525) \rightarrow \gamma\gamma}^{(th)} = \frac{12N_c}{5} Q_s^4 \alpha^2 \frac{|\mathcal{R}'_P(0)|^2}{m^4}, \quad (2)$$

where $N_c=3$ is the color number, $\alpha=\frac{1}{137}$, and fitting them with their experimental values 2.6 KeV and 0.081 KeV for $f_2(1270)$ and $f'_2(1525)$ respectively [3], we get

$$|\mathcal{R}'_{f_2(1270)}(0)|^2 = 6.6 \times 10^{-4} \text{ GeV}^5, \quad |\mathcal{R}'_{f'_2(1525)}(0)|^2 = 1.1 \times 10^{-3} \text{ GeV}^5. \quad (3)$$




FIG. 1: Feynman diagrams for charmonium ($^3S_1$) radiative decays into light mesons (^3P_J)

process	$J/\psi \rightarrow \gamma f_0(980)$	$J/\psi \rightarrow \gamma f_1(1285)$	$J/\psi \rightarrow \gamma f_2(1270)$	$J/\psi \rightarrow \gamma f'_1(1420)$	$J/\psi \rightarrow \gamma f'_2(1525)$
BR_{th}	1.6×10^{-4}	7.0×10^{-4}	8.7×10^{-4}	1.8×10^{-4}	2.0×10^{-4}
BR_{ex}	\	$(6.1 \pm 0.8) \times 10^{-4}$	$(13.8 \pm 1.4) \times 10^{-4}$	$(7.9 \pm 1.3) \times 10^{-4}$	$(4.5^{+0.7}_{-0.4}) \times 10^{-4}$

TABLE I: Numerical results for $\mathcal{B}(J/\psi \rightarrow f_J \gamma)$.

For charmonium, the value of the wave function at the origin is taken to be [10]:

$$|\mathcal{R}_{J/\psi}(0)|^2 = 0.81 \text{ GeV}^3, \quad |\mathcal{R}'_{\chi_{cJ}}(0)|^2 = 0.075 \text{ GeV}^5. \quad (4)$$

In the calculation, the meson masses are taken to be $M = 2m_q$ in the nonrelativistic limit. For the J/ψ decay we take $\alpha_s(2m_c) = 0.26$. Our results are listed in the Table I.

Then we will give the contributions of different helicities. As in Ref.[11], we choose the moving direction of f_J as the z-axis, and introduce three polarization vectors:

$$\omega^\mu(1) = \frac{-1}{\sqrt{2}}(0, 1, i, 0), \quad \omega^\mu(-1) = \frac{1}{\sqrt{2}}(0, 1, -i, 0), \quad \omega^\mu(0) = \frac{1}{m}(|\mathbf{k}|, 0, 0, k^0), \quad (5)$$

thus we can characterize the polarization tensor $\epsilon^{\mu\nu}(\lambda)$ of f_2 . With the same parameters, we give the branching ratios for different helicity states in Table II. The corresponding values of helicity parameters x and y are $x = 0.46$, $y = 0.23$. Recently, new experimental data for the contributions of different helicities in process $J/\psi \rightarrow \gamma f_2(1270)$ have been given by the BES Collaboration [12]: $x = 0.89 \pm 0.02 \pm 0.10$ and $y = 0.46 \pm 0.02 \pm 0.17$ (see also[3]). It is about 2 times larger than our results. But if we use a larger constituent quark mass, e.g. $m_u = M(1270)/2$, we will get substantially increased values $x = 0.79$ and $y = 0.58$ (also see Ref.[1]). We emphasize that the helicity parameters are very sensitive to the light quark masses, and hence very useful in clarifying the decay mechanisms. Note that if $m_u/m_c \rightarrow 0$, we will have $x \rightarrow 0$ and $y \rightarrow 0$, but this is inconsistent with data.

From Table I we see BR of $J/\psi \rightarrow f_0(980)\gamma$ (assuming $f_0(980)$ to be $s\bar{s}$) is predicted to be 1.6×10^{-4} . Actually, this calculation can also apply for $f_0(1710)$ (if treated as $s\bar{s}$) with an even smaller BR due to the phase space correction. However, experimentally BR of $J/\psi \rightarrow f_0(1710)\gamma$ is larger than 9×10^{-4} [3]. This implies that the $f_0(1710)$ can not be a pure $s\bar{s}$ or $(u\bar{u} + d\bar{d})/\sqrt{2}$ scalar meson (for the latter the BR is predicted to be about 4×10^{-4}). One possible explanation is that the $f_0(1710)$ has a substantial gluonic component due to $q\bar{q}$ -glueball mixing.

Next, we discuss J/ψ radiative decays to η and η' . Assume η' and η can be written as admixtures of flavor-singlet ψ_1 and octet ψ_8 :

$$\psi_1 = \frac{1}{\sqrt{3}}(u\bar{u} + d\bar{d} + s\bar{s}), \quad \psi_8 = \frac{1}{\sqrt{6}}(u\bar{u} + d\bar{d} - 2s\bar{s}), \quad (6)$$

$$\eta = \psi_8 \cos \theta - \psi_1 \sin \theta, \quad \eta' = \psi_8 \sin \theta + \psi_1 \cos \theta, \quad (7)$$

process	$ \lambda $	BR	process	$ \lambda $	BR
$J/\psi \rightarrow \gamma f_1(1285)$	1	0.18×10^{-4}	$J/\psi \rightarrow \gamma f'_1(1420)$	1	0.12×10^{-4}
	0	6.81×10^{-4}		0	1.64×10^{-4}
$J/\psi \rightarrow \gamma f_2(1270)$	2	0.38×10^{-4}	$J/\psi \rightarrow \gamma f'_2(1525)$	2	0.21×10^{-4}
	1	1.44×10^{-4}		1	0.51×10^{-4}
	0	6.91×10^{-4}		0	1.27×10^{-4}

TABLE II: Results for $J/\psi \rightarrow f_J \gamma$ with different helicity states.

process	BR_{th}	BR_{ex}
$J/\psi \rightarrow \gamma\eta$	3.0×10^{-4}	$(8.6 \pm 0.8) \times 10^{-4}$
$J/\psi \rightarrow \gamma\eta'$	2.3×10^{-3}	$(4.31 \pm 0.3) \times 10^{-3}$

TABLE III: Branching ratios for $J/\psi \rightarrow \eta(\eta')\gamma$.

process	BR_{th}	process	BR_{th}	process	BR_{th}
$\chi_{c2} \rightarrow \gamma\rho$	4.4×10^{-6}	$\chi_{c2} \rightarrow \gamma\omega$	5.0×10^{-7}	$\chi_{c2} \rightarrow \gamma\phi$	1.1×10^{-6}
$\chi_{c1} \rightarrow \gamma\rho$	1.4×10^{-5}	$\chi_{c1} \rightarrow \gamma\omega$	1.6×10^{-6}	$\chi_{c1} \rightarrow \gamma\phi$	3.6×10^{-6}
$\chi_{c0} \rightarrow \gamma\rho$	1.2×10^{-6}	$\chi_{c0} \rightarrow \gamma\omega$	1.3×10^{-7}	$\chi_{c0} \rightarrow \gamma\phi$	4.6×10^{-7}

TABLE IV: Predicted branching ratios for $\chi_{cJ} \rightarrow \gamma\rho(\omega, \phi)$.

where θ is the mixing angel, taken to be a widely accepted value $\theta = -20^\circ$ [13]. For the $\eta(\eta')$, the wave function at the origin may be determined from the leptonic decay of vector mesons $V \rightarrow e^+e^-$ ($V = \phi, \rho$), with the assumption that in the nonrelativistic quark model the wave functions of the 0^{-+} and 1^{-+} mesons should be equal in the nonrelativistic limit. Using

$$\Gamma(\phi(1020) \rightarrow e^+e^-) = N_c Q_s^2 \alpha^2 \frac{|\mathcal{R}(0)|^2}{3m_s^2} = (1.27 \pm 0.04) \text{ KeV}, \quad (8)$$

we can get $|\mathcal{R}(0)|^2 = 0.054 \text{ GeV}^3$ for $s\bar{s}$. Similarly we can get $|\mathcal{R}(0)|^2 = 0.032 \text{ GeV}^3$ for $u\bar{u}$ and $d\bar{d}$ from ρ leptonic decay. The decay branching ratios of $J/\psi \rightarrow \gamma\eta(\eta')$ are then calculated and listed in Table III.

We can see that for $J/\psi \rightarrow \gamma\eta(\eta')$ the simple nonrelativistic quark model calculations fit the data rather well, and the values are compatible with other approaches e.g. the $\eta(\eta') - \eta_c$ mixing mechanism based on the QCD axial anomaly[14].

Another interesting processes are decays of P-wave charmonia $\chi_{cJ} \rightarrow \rho(\omega, \phi)\gamma$ ($J=0,1,2$), where the photon is emitted by the light quarks but not the charm quarks, in contrast to the above discussed J/ψ radiative decays. The calculated branching ratios are shown in Table IV. In particular, the branching ratio of $\chi_{c1} \rightarrow \rho\gamma$ is predicted to be 1.4×10^{-5} , which may be tested by CLEO-c and BESIII in future experiments.

In this letter, we apply perturbative QCD to the radiative decays of J/ψ and χ_{cJ} into light mesons by treating the light mesons such as $f_2(1270)$ as nonrelativistic $q\bar{q}$ states as in the nonrelativistic constituent quark models. We perform a complete numerical calculation for the quark-gluon loop diagrams, where the light quark masses are taken to be the constituent quark masses. The calculated J/ψ decay branching ratios to the P-wave mesons $f_2(1270)$ and $f_1(1285)$ fit the data well (but the observed helicity parameter x for $f_2(1270)$ favors even larger light quark masses). Moreover, the results for S-wave mesons $\eta(\eta')$ seem to be also rather good. Our calculation does not support $f_0(1710)$ being the $s\bar{s}$ or $(u\bar{u} + d\bar{d})/\sqrt{2}$ meson, while the possible $s\bar{s}$ interpretation for $f_0(980)$ can be tested. The predicted branching ratios for $\chi_{cJ} \rightarrow \rho(\omega, \phi)\gamma$ ($J=0,1,2$) are generally small, and may be tested by CLEO-c and BESIII with future experiments.

However, we must bear in mind that for J/ψ decays the applicability of perturbative QCD is marginal, and the nonperturbative effects should be important. Moreover, the successes of nonrelativistic constituent quark models for light hadrons have not been fully understood from QCD. The constituent quark model description for η and η' is even more questionable. In this paper, to evaluate the quark-gluon loop diagrams and to avoid the soft and collinear divergences caused by massless light quarks, we have tried to use the constituent quark masses as in the nonrelativistic quark model for light mesons. Hopefully, the good (if not accidental) fit of the calculation presented here with the experimental data can shed light on a deeper understanding of J/ψ decays in QCD, which evidently deserve further studies.

We thank Ce Meng for valuable discussions. This work was supported in part by the National Natural Science Foundation of China (No 10421503), and the Key Grant Project of Chinese Ministry of Education (No 305001).

-
- [1] Körner J G, Kühn J H and Schneider H 1983 *Phys. Lett. B* **120** 444; Körner J G, Kühn J H, Krammer M and Schneider H 1983 *Nucl. Phys. B* **229**115.
[2] Krammer M, 1978 *Phys. Lett. B* **74**, 361 ; Korner J G and Krammer M, 1983 *Z. Phys. C* **16**, 279 ;
[3] Eidelman S et al. (Particle Data Group) 2004 *Phys. Lett. B* **592** 1.
[4] Close F E, Farrar G R and Li Z P 1997 *Phys. Rev. D* **55** 5749.

- [5] Cho P and Leibovich A K 1996 *Phys. Rev. D* **53** 150; Ko P, Lee J and Song H S 1996 *Phys. Rev. D* **54** 4312; Liu K Y, He Z G, and Chao K T 2003 *Phys. Lett. B* **557** 45.
- [6] Kühn J H 1979 *Nucl. Phys. B* **157** 125; Guberina B et al 1980 *Nucl. Phys. B* **174** 317.
- [7] Mertig R, Böhm M and Denner A 1991 *Comput. Phys. Commun.* **64** 345.
- [8] Hahn T and Perez-Victoria M 1999 *Comput. Phys. Commun.* **118** 153.
- [9] Denner A and Dittmaier S 2003 *Nucl. Phys. B* **658** 175.
- [10] Eichten E J and Quigg C 1995 *Phys. Rev. D* **52** 1726.
- [11] Ma J P 2001 *Nucl. Phys. B* **605** 625; 2001 *Erratum – ibid.* B **611** 523.
- [12] Ablikim M et al hep-ex/0603048.
- [13] Gasser J and Leutwyler H 1985 *Nucl. Phys. B* **250** 465.
- [14] Chao K T 1989 *Nucl. Phys. B* **317** 597; Chao K T 1990 *Nucl. Phys. B* **335** 101.